

University of Queensland

THE EFFICACY AND COGNITIVE PROCESSES OF
COGNITIVE BEHAVIOUR THERAPY IN THE
TREATMENT OF PANIC DISORDER WITH
AGORAPHOBIA

Tian P.S. Oei, Michael Llamas, and Grant J. Devilly *

School of Psychology.
University of Queensland,
Brisbane. Queensland 4072
Australia.

Running head: PANIC DISORDER WITH AGORAPHOBIA

Acknowledgment: We would like to thank Dr L. Evans and staff at the Anxiety Disorders Clinic for their support and use of facilities.

* All authors contributed equally to this paper. Requests for reprints to Dr T.P.S. Oei.

ABSTRACT

Cognitive behaviour therapy (CBT) has been shown to be an effective therapy for a variety of clinical problems. The present paper aimed firstly to review the literature to examine the effectiveness of CBT as treatment for PDA. Additionally, demonstrations of the efficacy of CBT treatments have largely been accepted as evidence to confirm the cognitive models of psychopathology which propose that cognitive processes are primarily responsible for abnormal behaviour. To date, the evidence for the assumption that CBT supports cognitive models of PDA has not been systematically reviewed. The second aim of this paper, therefore, was to examine any such evidence. A literature review of CBT studies in the area of PDA was conducted using both descriptive and quantitative (meta-analysis) procedures. In all, thirty five studies published between 1969 and 1996 were included in this review. The results show that CBT is an effective treatment for PDA. However, the contribution of cognitive processes to this disorder and the role that they play in the successful outcome of CBT remain unclear and in need of further empirical investigation. At present, CBT provides little support to validate the cognitive models of PDA.

Cognitive behaviour therapy (CBT) is becoming increasingly more popular in the treatment of anxiety disorders. This marked growth of interest in the use of CBT with anxiety disorders has, to a large extent, been due to an increasing awareness of the importance of cognitive processes in these and other emotional disorders (Emmelkamp, 1982; Free & Oei, 1989; Oei, Duckham & Free, 1989; Oei, Lim & Young, 1991; Powell & Oei, 1991; Oei & Free, 1995; Oei & Shuttlewood, 1996). One anxiety disorder which has received particular attention has been agoraphobia with panic attacks (APA, 1980) or panic disorder with agoraphobia (PDA)(APA, 1987; 1994). This condition refers to a well recognised clinical syndrome encompassing panic attacks, phobic anxiety and avoidance, as well as anticipatory anxiety or fear of panic. The central notion of CBT is the idea that cognitions and information processing, in particular catastrophic cognitions, are central in the causation and maintenance of PDA (Clark, 1988; Lang, 1988; Rapee, 1987; Salkovskis, 1988; Khawaja & Oei, in press). Consequently, many consider that therapy should involve the utilisation of cognitive techniques to modify maladaptive catastrophic cognitions and to produce psychological and behavioural changes, while others may view any intervention as being able to create cognitive shifts (see Chambless & Gillis, 1993, for review).

Two issues which require further attention relate to the efficacy and underlying processes (or assumptions) of CBT in the treatment of PDA. While various reviews have examined the efficacy of CBT in the treatment of depression (Oei & Free, 1995); social phobia (Heimberg, 1989) and alcohol dependence (Oei et al., 1989) there are currently few extensive reviews of the relevance of CBT for panic disorder without agoraphobia (e.g. Beamish, Granello, Granello, McSteen, et-al, 1996) and none for PDA. This is surprising given that PDA is one of the most debilitating conditions amongst the anxiety disorders (see Weissman, 1990, 1991; Weissman & Merikangas, 1986). Consequently, the present review will firstly seek to examine if CBT is an efficacious treatment for PDA.

In addition, several cognitive models of causative processes have been highlighted in the literature in relation to PDA (see Rachman & Maser, 1988). Beck (1988), for example, claims that PDA is precipitated by danger signals which are based on the interpretation of internal sensations as indicative of heart attack and dying. Similarly, Clark (1988) and Rapee (1986, 1987) propose that panic disorder originates from the catastrophic interpretations of certain bodily sensations; for example, perceiving palpitations as evidence of impending heart attack or racing thoughts as evidence of going crazy. Additionally, Lang (1988) views PDA in the context of a faulty information processing system designed to control the pattern, sequence and timing of behaviour. However, irrespective of how one conceptualises the role of cognitions in PDA, adherents of cognitive theories of this disorder commonly hold cognitive variables to be of primary importance. As a result, the assumption has implicitly or explicitly been made that cognitions cause and maintain PDA. Consequently, there is a belief among

clinicians that demonstrations of the efficacy of CBT treatments for PDA will provide the sufficient evidence required to confirm the cognitive models of such psychopathology. Such an inference has been argued by Oei, et al. (1989) to be invalid. To date, there is no systematic review of empirical evidence to support this assumption in the literature. The second aim of the present paper, therefore, is to examine if the CBT literature supports the cognitive models of PDA.

CBT AND PANIC DISORDER WITH AGORAPHOBIA: OVERVIEW OF STUDIES

Two computer searches of the literature on CBT and PDA was conducted to identify the relevant papers that have researched this area during from the period 1969 to 1989 and from 1990 to 1996. Two major data bases, PsycLit CD and Carl online, were searched, using the keywords panic disorder, agoraphobia, treatment and outcome. On the search from 1990 to 1996, 45 papers were highlighted. Studies employing CBT were included in the review if the CBT had been applied to subjects who met the criteria for agoraphobia with panic attacks (APA, 1980), or PDA (APA, 1987; 1994), or who would have met either of these criteria had they been applied. Studies were excluded if they were a sub-set of an already included study, a single case study, a review of the literature, or written in a language other than English.

In all, thirty five empirical studies were identified which fulfilled the above criteria. However, this is not to infer that the current review is exhaustive, indeed considering the rapid rate of research within this area, even if this were exhaustive, by the time of going to press this status would have changed. Rather this paper aims to provide an indication of the trend to non-researchers who may have neither the time nor the resources to investigate this particular area in any depth and to produce an estimate of the research climate so that our comments can be appreciated within this backdrop. Table 1 summarises these studies.

Insert Table 1 about here

It can be seen from table 1 that a number of research groups have been involved in conducting studies which meet the criteria outlined above. In total, 1317 patients have been studied. Of the studies which delineated between gender, 761 out of 948 (80.27%) patients were female. This figure is consistent with reports in the literature that PDA is more frequently diagnosed in women (eg. Oei, Wanstall & Evans, 1990).

Several research designs have been employed by the studies listed in table 1 including: a) group comparison (eg. Ascher et al., 1986); b) single case-study (eg. DeVoge et al., 1981); c) multiple baseline across subjects (eg. Last et al., 1984); d) Post Hoc design (Chambless & Williams, 1995);

e) cross-over (eg. Emmelkamp et al. 1978); f) sequential staggering of treatment conditions (e.g., Ascher, 1981) and g) time series and repeated measures (eg. Salkovskis et al., 1986). Seventeen studies used randomization as part of their research design and only four studies included a clean control group.

A number of CBT approaches are represented by the studies in table 1 including: cognitive restructuring and training, self-statement training, paradoxical intervention, covert rehearsal of coping with anxiety, reattribution of somatic symptoms, coping thoughts training, thought stopping, breathing and relaxation training and variations of prolonged exposure. Twenty studies investigated the effects of CBT on PDA but did not include other treatment comparison groups in their evaluations. The remaining studies, however, did attempt to compare the CBT to other forms of therapy (e.g., exposure alone, fluvoxamine, etc) or to varying versions of CBT itself.

With regard to outcome measures, eight studies employed few such measures while the remainder tended to employ large and comprehensive assessment batteries. Fewer still included physiological as well as cognitive and behavioural assessments as part of their outcome measures. Thus there are only a limited number of studies available that take into account the tripartite (cognitive, behavioural and physiological) nature of PDA in their assessments. Furthermore, there is a large variability and lack of agreement that exists amongst researchers regarding the assessment techniques and instruments used when researching PDA. Clearly, more agreement and consistency in assessment strategies needs to exist if comparisons of results across future studies is to be made easier. Additionally, future researchers need to take more account of the "triple response" nature (Himadi, Boice & Barlow, 1985) of PDA in their assessments.

Finally, table 1 also shows that most studies, with the exception of eight (Ascher, 1981; Ascher et al., 1986; Barlow et al., 1984; Marchione et al., 1987; Michelson et al., 1990; DeBeurs et al., 1993; 1995; Hoffart, 1995), included follow-up periods as part of their investigations. These reported follow-up periods ranged between 1 to 16 months and one study of 9 years.

THE EFFICACY OF CBT IN THE TREATMENT OF PANIC DISORDER WITH AGORAPHOBIA

In an attempt to answer the question of whether CBT is efficacious in the treatment of PDA, the present section will aim to analyse the available literature in two ways. Firstly, the different types of outcome measures that have been employed by the studies in table 1 will be reviewed with a view to determining if CBT was associated with any significant change in these. Secondly, a meta-analysis of those studies in table 1 reporting data on the Fear Questionnaire will be conducted. To the extent that CBT is an effective treatment of PDA, one would expect the results of both of these analyses to converge towards supporting such a conclusion.

ANALYSIS OF OUTCOME MEASURES

A number of outcome measures (domains) were employed by the studies in table 1. These will now be discussed in terms of whether CBT was associated with any significant change in each of them or not. Table 2 lists the results of the studies that report outcome data on the various domains.

 Insert Table 2 about here

1) Panic Measures

Of the 35 studies reported in table 1, 60% reported outcome from measures of panic. Of these 100% reported improvement by post-treatment and /or follow-up (see table 2). Hence CBT has consistently been associated in the PDA literature with significant and positive changes on a variety of panic measures both at post-treatment and for up to 6 months follow-up, and in one study, 9 years follow-up.

2) Fear and Avoidance Measures

A host of measures of fear and avoidance have been utilised in the assessment of PDA, including the Fear Questionnaire, the Mobility Inventory, and the Phobic Anxiety and Avoidance Scale. Seventy seven percent of the studies reviewed utilised measures of fear and avoidance, and of these, 100% reported improvement by post-treatment and /or follow-up. Overall, it can therefore be said that CBT treatment of PDA has been consistently associated in the literature with significant post-treatment and follow-up improvements of up to 16 months on fear and avoidance as measured by most of the major assessment instruments employed in this area.

3) Severity/Intensity Measures

Measures of severity and intensity have most commonly included SUD scales and the Global Assessment of Severity Scale, amongst others. Forty six percent of the studies reviewed reported outcome for this domain, of which 100% reported improvement by post-treatment and / or follow-up. Of those studies that included follow-up data only the Mavissakalian et al. (1983) study did not find improvements during post treatment to follow-up analysis. With the exception of this one study, CBT therefore appears to be associated with improved changes in severity and intensity measures of PDA both at post treatment and follow-up periods ranging up to 16 months.

4) General Anxiety Measures

Fifty four percent of the studies utilised measures of general anxiety, of which 90% report statistical improvement. Two studies (Marchione et al., 1987, and Emmelkamp et al., 1978) did not find statistical improvement. In the Marchione et al (1987) study perusal of the data by the authors showed that CBT did produce substantive therapeutic gains pre- to post-treatment on the Taylor

Manifest Anxiety Scale, however these changes did not achieve conventional levels of statistical significance. The reason for this was, in part, attributed by the authors to the small sample size used in their study. Also, with regards to anxious mood, Emmelkamp et al. (1978) initially found that CBT which included prolonged exposure as a treatment component of PDA was not associated with statistically significant improvements on the Anxiety Mood Scale at post-treatment. In a later study however, Emmelkamp and Mersch (1982) did find CBT to be associated with post-treatment improvements on the AMS irrespective of whether it incorporated a prolonged exposure component or not. In general therefore, apart from the studies by Emmelkamp et al. (1978) and Marchione et al. (1987), the literature to date does provide evidence that CBT treatment of PDA can lead to significant improvements on a variety of measures of general clinical anxiety. Furthermore, these improvements have been observed up to 6 months follow-up.

5) Social Anxiety Measures

Measures of social anxiety have included the Fear of Negative Events Questionnaire, the Social Avoidance and Distress Scale and the Social Phobia subscale of the Fear Questionnaire. Of the 35 studies cited, only 14% utilised a measure of social anxiety. Of these, 4 reported improvement and one reported no statistical effect. This study (Beck et al., 1994) did display a reduction in the cognitive therapy group on the social phobia subscale of the Fear Questionnaire, but this did not reach statistical significance. Overall, it would appear that CBT generally decreases social anxiety at post-treatment and that this is maintained up to 12 months follow-up.

6) General Symptomatology Measures

Twenty six percent of the studies cited utilised measures of general symptomatology (e.g. Hopkins Symptom Checklist, Subjective Symptoms Scale, Subjective Symptoms Checklist, Lehrer and Woolfolk Symptom Questionnaire, Symptom Checklist 90-R). Of these, all reported improvement to post-treatment and / or follow-up, with at least one of these measures. However, Marchione et al. (1987) report inconsistent results, displaying improvement on the Subjective Symptoms Checklist but not on the Hopkins Symptom Checklist. Except for this study CBT was associated with improvement on general symptomatology which was maintained to 6 month follow-up.

7) Behavioural Measures

As can be seen from table 2, 46% of the studies reported outcomes for behavioural measures, of which only one study (Last et al., 1984) reported mixed results and another (Mavissakalian et al., 1983) found no significant improvements. Hence, whilst there are some inconsistencies noted amongst studies to date concerning the impact of CBT used in PDA on behavioural measures, the majority have shown CBT to be associated with significant improvements on these measures both at post-treatment and at follow-up periods ranging up to 12 months.

8) Cognitive Measures

Only 46% of the studies reported using cognitive measures, of which all except one (Last et al., 1984) reported positive results. Further discussion of outcome associated with CBT on these measures will be postponed till a later section of the paper as they are relevant to the question of whether CBT supports cognitive models of PDA.

9) Physiological Measures

Of those studies in table 1, four (Last et al., 1984; Marchione et al., 1987; Michelson et al., 1985; Michelson et al., 1990) employed Heart Rate (HR) as a physiological outcome measure of PDA, and one (Beck et al., 1994) used HR, skin conductance and trapezius electromyographs(EMG). Results of these studies are somewhat inconsistent. In the Marchione et al. (1987) study of agoraphobics and the Michelson et al. (1990) study, CBT was associated with significant pre to post-treatment improvements observed on most HR measures taken during a behavioural avoidance test. Michelson et al (1985), in contrast, found that patients with PDA who underwent CBT actually experienced increases in physiological reactivity during treatment, and it was not until 3 months follow-up that significant improvements were noted for CBT on HR sitting and HR walking measures. In the Beck et al. (1994) study there was found to be no significant effects for HR and EMG, and conflicting statistical results by treatment group for skin conductance. The authors attribute this to pre-treatment differences between groups on this measure. Finally, Last et al. (1984), reporting data on two agoraphobic patients who underwent CBT, found that the magnitude of the HR measure for one patient was higher during post-treatment assessment, whereas for the second patient it was much lower when compared to pre-treatment. Hence, studies that have to date evaluated the impact of CBT on physiological measures of PDA are very limited in number, have focussed primarily on heart rate responses and have led to somewhat inconsistent findings.

10) Locus of Control

Seven studies (20%) from table 1 included Locus of Control Scales in their assessment batteries. Statistically significant improvements in these scores were reported with CBT interventions across treatment and follow-up phases of up to 6 months. Hence, taken together, the results of these studies indicate that CBT can be associated with significant and positive changes in locus of control when used as a treatment of PDA.

11) Depression Measures

Sixty six percent of the studies in table 1 measured depression, and by far the most commonly employed measure was the Beck Depression Inventory. All of the studies reported a significant reduction in depression. Overall, when one considers this literature in relation to PDA, it can be concluded that CBT has consistently been associated with reductions in depression as measured by the

major scales currently employed by researchers. Further, these reductions have been evidenced at post-treatment and follow-up phases of up to 12 months.

12) Marital State Measures

It can be seen from table 1 that four studies (Barlow et al., 1981; Barlow et al., 1984; Chambless et al., 1986; Michelson et al., 1988) evaluating CBT with PDA have employed measures of marital state. Two studies (Barlow et al., 1984; Michelson et al., 1988) included the Marital Adjustment Scale in their assessment measures. Significant improvements were associated with CBT on this scale in the Michelson et al., (1988) study on both post and 3 month follow-up scores. In the Barlow et al (1984) study no specific results were mentioned for this scale, however, a statement made by the authors implied that improvements did occur pre and post treatment when CBT was employed.

Two studies (Barlow et al., 1981; Barlow et al., 1984) used the Marital Happiness Scale (MHS). Using a CBT treatment with agoraphobics, Barlow et al (1981) found that two patterns of relationship emerged with regards to marital happiness and severity of phobia. For four of the couples treated in their study, as phobia improved marital happiness increased. For two couples though, the inverse relationship was noted whereby improvement in phobia was correlated with decreases in marital happiness. No specific results were mentioned by the Barlow et al., (1984) study relating to the MHS, however, a statement made by these authors implied that improvements did occur pre to post treatment when CBT was similarly employed with agoraphobics.

Finally, Chambless et al. (1986) utilised the Marital Dissatisfaction Questionnaire (MDQ) in attempting to search for predictor variables of CBT treatment outcome for PDA. These authors found that patients with higher pre-treatment scores on the MDQ were no less likely to improve in their program than patients with lower pre-treatment scores on the MDQ. However, since post-treatment and follow-up measures were not obtained, it is not possible to say in this study whether the CBT may have had any impact on marital dissatisfaction scores across time.

Overall, the impression that one acquires from perusing this area of the literature is that CBT treatment of PDA does generally appear to impact on marital state, but that this may not always be in the positive and clinically therapeutic direction.

13) Assertiveness Measures

As can be seen from table 1, five studies (Chambless et al., 1986; Emmelkamp et al., 1978; Emmelkamp & Mersch, 1982; Michelson et al., 1985; Michelson et al., 1988) have considered the impact that CBT has on assertiveness as part of their investigations on PDA. Each of these studies, apart from Emmelkamp et al. (1978) found significant associations between a CBT intervention and improvements in assertiveness. Chambless et al. (1986) reported significantly improved change in their patients with PDA during a 6 month period following an intensive 2-week CBT treatment program on

the Gambrill and Rickey Assertiveness Inventory. Similarly, Emmelkamp & Mersch (1982) found significant changes on the Adult Self-Expression Scale following CBT treatment of PDA and at 1 month follow-up. In the Michelson et al. (1985; 1988) studies patients with PDA showed improvements on the Wolpe-Lazarus Assertiveness Inventory at post treatment and at 3 months follow-up. Needless to say, the studies that have employed assertiveness measures in their research of PDA are few in numbers. Nevertheless the results of those that have been conducted to date suggest that, on the whole, CBT can be associated with improved changes in assertiveness scores both at post-treatment and at follow-up periods of up to 6 months.

14) Endstate Functioning and Improvement

Table 1 shows that ten studies have used a-priori operationalized composite measures of Endstate Functioning and Improvement as indices of outcome for CBT treatment of PDA. These have included the Operationalised Measure of Endstate Functioning, the Operationalised Measure of Endstate Improvement and the Composite Measure. Operationalized definitions of endstate functioning and improvement have the merit of combining several measures in different response systems to produce a more composite score on which to base treatment outcome. Using this approach, the classification of high endstate functioning and improvement would represent superior clinical status with negligible or no phobic symptoms. From table 2 one can see that CBT has been consistently associated with, at minimum, moderate numbers of patients with PDA acquiring high endstate functioning and improvement when such measures have been employed following treatment and at follow-up periods of up to 6 months.

15) Other Outcome Measures

In addition to those outcome measures already discussed above, several studies from table 1 have reported using measures such as Valium Intake (VI), Car Mileage (CM), Daily Self Appraisal Interviews (DSAI) and Rate of Hospital Admissions (RHA) (DeVoge et al., 1981) as well as Biographical Data Sheets (BDS) (Chambless et al., 1986), Home Visit Observations (HV) (Barlow et al., 1984) and Behavioural Diaries (BD) (Barlow et al., 1984; Michelson et al., 1986). It is not possible to state the impact of CBT on the BDS, HV and BD due to the nature of these measures and the way they were reported in their respective studies. The results on the measures used by DeVoge et al. (1981) (VI, CM, DSAI and RHA) however are noteworthy. That is, CBT used to treat a severe agoraphobic patient was found to produce desirable changes in VI, DSAI and CM (or avoidant behaviour). Furthermore, these changes were accompanied by fewer hospitalisations of the patient during a 16 month follow-up period.

Summary: To summarise therefore, the above discussion would suggest that CBT does lead to positive and concurrent changes on several measures that have been employed by available studies

in the area of PDA. Specifically, CBT when used as a treatment of PDA has been associated with significant and positive changes on measures of panic, fear and avoidance, approach behaviours, severity and intensity of condition, endstate functioning and improvement, clinical anxiety, depression, assertiveness and locus of control. CBT has also been associated with changes in marital state, however these changes have not always been in the positive and clinically therapeutic direction. Furthermore, the noted changes on the measures described have been shown to occur both at post-treatment and at follow-up periods ranging up to 16 months. Finally, the above discussion also highlights several measures that have been employed in the literature to date on which the impact of CBT, when employed with PDA, still remains unclear.

ANALYSIS 3: META-ANALYSIS OF STUDIES IN TABLE 1

An alternative method of determining the effectiveness of an intervention has been suggested by Nietzel and Trull (1988). This method provides a useful tool for quantifying the clinical significance of treatment outcomes by comparing the pre, post and follow-up scores with normative scores obtained from “normal” individuals (Trull, Nietzel and Main, 1988; Nietzel and Trull, 1988). For the purposes of our meta-analysis, normative data previously reported by Nietzel and colleagues for the Fear Questionnaire (FQ), for both a college group and community sample, was employed. This was due to the FQ’s reportedly frequent use in the agoraphobia outcome literature, it’s demonstrated psychometric properties and its endorsement by a number of leading agoraphobia researchers. In order to compare the results of clinical outcome studies with the normative distributions, a standardised effect size score is computed for the relevant treatment group in each study using the formula $(M_t - M_{n1}) / SD_{n1}$ and $(M_t - M_{n2}) / SD_{n2}$, where M_t is the mean post-treatment Agoraphobia or Total Phobia subscale score of the FQ; M_{n1} and M_{n2} are the mean FQ norms for the college and community samples respectively; and SD_{n1} and SD_{n2} are the corresponding standard deviations for these two normative samples. Similarly, effect sizes that contrast the normative groups with treated groups pre-test and follow-up scores are also computed.

Of the studies identified in table 1, eleven reported the FQ agoraphobia subscale scores in their text, and seven reported total FQ scores. Effect sizes for the CBT group in these studies were calculated for pre-test, post-test and follow-up results on the FQ using the norms of Nietzel and Trull (1988). These are reported in table 3 (for the Agoraphobia subscale) and in table 4 (for the Total Phobia subscale).

 Insert Tables 3 and 4 about here

Mean effect sizes of studies reported in table 3 were calculated for pre, post and follow-up phases of treatment (controls not included). These are shown in figure 1 plotted against the distributions for the normative groups obtained by Nietzel and Trull (1988) for the FQ Agoraphobia subscale.

 Insert Figure 1 about here

It can be seen from figure 1 that, on the FQ Agoraphobia subscale, patients on the average, moved from 3.88 to 1.70 standard deviations of the collegiate mean at post-treatment and to 1.70 standard deviations at follow-up. They also moved from 1.37 to 0.17 standard deviations of the general population mean at post-treatment and to 0.24 standard deviations at follow-up. Hence, CBT was associated with improvements on FQ Agoraphobia scores from pre to post-treatment and follow-up as demonstrated in both analyses. Jacobson, Follette and Revensdorf (1984) suggest that scores within two standard deviations of the “normative” groups be considered clinically significant. It can therefore be concluded (based on figure 1) that the CBT treatment resulted in post-treatment and follow-up scores which are substantially clinically significant, and similar to FQ Agoraphobia subscores found in the general (community) population. In contrast, the CBT treatment results were only marginally clinically significant when compared to FQ Agoraphobia subscores found in a college population.

The mean effect sizes of studies reported in table 4 were calculated and are shown in figure 2, plotted against the distributions for the normative groups obtained by Nietzel and Trull (1988) for the FQ Total subscale.

 Insert Figure 2 about here

It can be seen from figure 2 that, on the FQ Total subscale, patients on the average moved from 2.11 to 0.38 standard deviations of the collegiate mean at post-treatment and to 0.29 standard deviations at follow-up. They also moved from 0.97 to -0.48 standard deviations of the general population mean at post-treatment and to -0.47 standard deviations at follow-up. It can therefore be concluded (based on figure 2) that the CBT treatment resulted in improved post-treatment and follow-up scores which are clinically significant and similar to FQ Total scores found in college and general populations. Further, these improvements were well maintained at follow-up.

When comparing figure 1 with figure 2, it further becomes apparent that the efficacy of CBT treatment is more pronouncedly demonstrated when the FQ Total subscale (as opposed to the FQ Agoraphobia subscale) is used as a measure of change. One possible explanation for this is that while CBT may be producing some measurable change on the FQ Agoraphobia subscale, it may also be having an impact on the Social Phobia and Blood/Injury Phobia ratings of this questionnaire. Various authors have either highlighted or alluded to the important role that catastrophic cognitions also play in social phobia (e.g., Arnkoff & Glass, 1989; Buttler, 1989) and in blood and injury phobias (Rachman & Maser, 1988). Hence, one would expect CBT to be effective in modifying catastrophic cognitions associated with these two phobias as well as with PDA. This being the case, the greater effectiveness noted with CBT on the FQ Total measure (compare figures 1 and 2) may well be reflecting the additive impact that this treatment may have had on the social and blood/injury phobia subscales that make up the FQ Total score. Significant changes in FQ Social and Blood/Injury subscores were in fact reported by some of the studies (e.g., Michelson et al., 1988) employed in the meta-analysis conducted by the present authors. Hence, there is some support for the hypothesis that CBT, when employed in the treatment of PDA, may not only be effecting agoraphobic catastrophic cognitions but may also indirectly be having an impact on social and blood/injury catastrophic cognitions at the same time.

Summary: The above results would suggest that CBT used in the treatment of PDA is effective in reducing self-reported agoraphobic symptoms as measured by the FQ. Furthermore, it would appear that the extent of clinical efficacy demonstrated by CBT when using this outcome measure will vary depending on the type of comparison norms that are used and the nature of the FQ subscales being utilised. Using Jacobson et al.'s (1984) criteria of falling within two standard deviations of a normal population's mean, the clinical efficacy of CBT with PDA was substantially demonstrated with community and collegiate norms, as reference groups at post-treatment and follow-up on the FQ Total scale. However, on the FQ Agoraphobia subscale, this criterion was substantially satisfied with the community norms but only marginally so with the collegiate norms. Additionally, the clinical efficacy of CBT was more pronouncedly noted on the FQ Total scale than on the FQ Agoraphobia subscale.

Overall Summary and Conclusion: The present section of this paper has sought to determine if CBT is efficacious in the treatment of PDA. Two analyses were conducted on the literature available to date. Firstly, the literature was reviewed specifically in terms of the association between CBT and the different outcome measures that have been employed. Secondly, a meta-analysis was conducted on the available studies which utilised the Fear Questionnaire as an outcome measure. In general, the results of these two analyses would seem to converge and lead to the conclusion that CBT is an effective treatment of PDA at least in the short-term. The next section of the paper will briefly discuss the current status with regards to the long-term efficacy of CBT with PDA.

LONG-TERM EFFICACY OF CBT WITH PANIC DISORDER WITH AGORAPHOBIA

Twenty six of the studies from table 1 included follow-up assessments. These ranged from 1 to 16 months after termination of treatment, and one study of 9 years follow-up. The most frequently used length of follow-up was 6 months. Thus it would be difficult to make any definitive conclusions about the long-term maintenance of effectiveness of CBT in treating PDA. There is a need for future research studies investigating the long-term outcome of CBT at lengthy follow-up intervals. The one study, in table 1, to investigate long-term follow-up treated with exposure in vivo only (Fava et al., 1995), yet found 67.4% of the those treated to continue remission for over 7 years. Closer inspection of the results of the Emmelkamp & Mersch (1982) study is interesting also in that they suggest that CBT efficacy may become more obvious over extended periods of assessment rather than during shorter periods. This is evidenced in their study of PDA patients by the way that exposure in vivo and a combined procedure (Self-statement training plus exposure in vivo) were superior to cognitive restructuring at post-test but at 1 month follow-up the differences in the treatments disappeared due to continuing improvement in the cognitive restructuring condition. In a similar manner the study by Mavissakalian et al. (1983) also provides evidence suggesting that Self-statement training requires a similar 'catching-up' period when compared to a Paradoxical intervention. Hence, in addition to the research needed to determine the long-term efficacy of CBT with PDA, research is also necessary to ascertain the minimum period required for CBT to demonstrate its effective results in the treatment of PDA.

DOES CBT SUPPORT COGNITIVE MODELS OF PANIC DISORDER WITH AGORAPHOBIA?

Demonstrations of CBT's efficacy are insufficient to infer a support for the cognitive models and processes of PDA. To do this at least three things must also be shown. Firstly, it must be shown that CBT is not only effective but that it also leads to change in the person's cognitions. Secondly, it needs to be demonstrated that effective CBT uniquely produces cognitive change in comparison to other therapies. Thirdly, it needs to be demonstrated that it is the change in cognitions that has produced any demonstrated improvement.

Table 1 shows that 15 out of 16 studies which included an independent measure of cognition met the first criterion of demonstrating the efficacy of CBT with PDA and linking this to a significant change in the cognitive measure. The remaining studies reviewed in table 1 did not include an independent measure of cognition. Mackay and Liddell (1986), although they did incorporate a 'cognitive' measure of self-efficacy into their assessment battery, did not comment sufficiently on its results in their text to enable useful conclusions to be drawn. Whilst the study by Last et al. (1984) did

not demonstrate clear and consistent changes in cognitions, the other studies which employed cognitive measures (see table 2) did demonstrate such changes. For example, in the Marchione et al. (1987) study, patients with PDA were instructed to verbalise their internal dialogue during a behavioural test. These verbalizations were subsequently classified and analysed in terms of whether they were self-defeating statements, coping statements or neutral/task-irrelevant statements. CBT was found to be associated with significant reductions in the percentage of negative statements and with significant increases in the frequency of positive statements from pre to post-treatment. Such a trend was mostly found in the other studies too.

It must be noted, however, that changes in cognitive measures employing tape-recorded verbalizations and/or thought lists may be due to experimental artefacts such as therapist reinforcement (see for example, Oei et al., 1989). In addition, Oei et al. (1989) argue that the cognitions to be measured should be the theoretical constructs that are postulated to influence the abnormal behaviour. In PDA the cognitions must be of a catastrophic nature. Chambless et al. (1986) did in fact attempt to measure changes in such cognitions related to PDA using the Agoraphobic Cognitions Questionnaire and the Body Sensations Questionnaire. Both of these questionnaires are designed to measure 'fear of fear' among agoraphobics (Chambless et al., 1984), a concept that, when linked with catastrophic cognitions is argued to mediate the escalation of anxiety to panic as occurs in PDA (Clark, 1988). In their study, Chambless et al. (1986) demonstrated significant changes on both the Agoraphobic Cognitions and Body Sensations questionnaires following successful CBT intervention with their panic disordered patients. In doing so, these authors provide strong supportive evidence for a link between CBT and a cognitive processes model of PDA.

In terms of the second criterion required in support of a cognitive processes model of PDA, effective CBT must be shown to produce cognitive change unique from other therapies. Two studies of PDA reviewed in table 1 (Marchione et al., 1987; Williams & Rappoport, 1983) have in fact demonstrated that, in addition to CBT, behavioural therapies (such as exposure) on their own, or in combination with other procedures, can also produce cognitive changes. Such results have led some authors (e.g., Williams & Rappoport, 1983) to propose that cognitive changes more likely occur following behavioural change. Authors such as Williams and Rappoport (1983) discuss this in terms of Bandura's (1977) self-efficacy theory which holds that fear is rooted in thought, but that the means by which thought is changed is through performance-based treatments. Clearly this issue requires more clarification through further research. It would also be useful to determine if the inclusion of a cognitive component to the behavioural therapy would serve as having an additive effect on the extent of cognitive change produced. In actual fact Marchione et al. (1987) and Williams and Rappoport (1983), to mention but two, did compare the addition of a cognitive component along with behavioural therapy

to the effects of behavioural therapy alone. In both of these studies the combined cognitive and behavioural procedure was found to be superior to the behavioural procedure alone in terms of changing positive coping thoughts but not fearful thoughts. Thus while other therapies such as exposure do appear to produce cognitive changes on their own, the CBT appears to **add** to these effects. This conclusion, however, is based on the results of only a few studies and thus should be accepted only with some caution at this stage.

Finally, the third criterion for a cognitive processes model of PDA requires that cognitive change be the active ingredient that leads to any improvement in the disorder. It is evident from the discussion thus far that in a minority of studies that have monitored cognitions, changes in these cognitions have in fact coincided with recovery from the PDA. This correlative link between cognitive change and recovery from PDA suggests some support for cognitive explanations of the disorder. This link has been further supported by more recent experimental studies, not included in this review, such as that by Salkovskis, Clark, Hackman, Wells & Gelder (in submission). However, it should be noted that other plausible aetiological factors need to be excluded before it can be solely concluded that cognitive change itself is the mechanism of action in CBT. For example, a study by Bandura, Taylor, Williams, Mefford and Bauchas (1985) suggests that the construct of self-efficacy may in fact interact with epinephrine and norepinephrine to produce fear reduction in patients with spider phobia. The implication of this study, when generalised to the condition of PDA, is that physiological variables may contribute an interactive role in the maintenance of and recovery from this condition. This suggestion is quite consistent with the proposed 'circular processes' model of depression by Free and Oei (1989) implicating both psychological and physiological aetiological contributions to the disorder. None of the studies reviewed in table 1 have evaluated physiological as well as psychological processes models of PDA in their research. It would therefore be impossible to conclude that any noted recovery in the PDA condition was due to cognitive change alone.

In summary, the question of whether CBT supports cognitive processes models of PDA still remains unclear. At this stage we can say that the efficacy of CBT has been well demonstrated and that there are a reasonable number of studies reviewed that have linked this to an independent measure of cognitive change. While this in itself provides some support for the cognitive processes models of PDA, the issue remains complicated in that CBT does not appear to produce cognitive change in isolation of other procedures such as exposure. Furthermore, the studies reviewed in the present paper have not adequately evaluated the likely contributions of other plausible aetiological and maintaining factors (such as biological processes) in their research of PDA.

CONCLUSION

The present review firstly sought to examine if CBT is an efficacious treatment for PDA. Secondly, it sought to determine if the CBT literature to date supports cognitive models of PDA. With respect to the first question, 35 empirical studies in the area were analysed descriptively and quantitatively by meta-analysis. The results of the analyses converged in support of the conclusion that CBT is therapeutically effective in the treatment of PDA. With regards to the second question, 16 (46%) of the empirical studies reviewed included cognitive measures in their assessments, of which 15 were able to link significant and positive changes on these measures to the CBT intervention employed to treat the PDA. However, some of the studies (e.g. Marchione et al., 1987; Williams & Rappoport, 1983) have shown positive change in the cognitive measures when behavioural therapies (such as exposure) have been used on their own or in combination to treat PDA. Consequently, the mechanisms by which CBT achieves its effects on this debilitating condition continue to remain somewhat unclear. While this review provides some evidence to suggest that cognitive processes may play an important role in the development and maintenance of PDA, they may not of themselves be sufficient to explain the condition. The empirical studies reviewed have not adequately evaluated how such cognitive processes may in fact be combining with and / or interacting with other mechanisms (such as biological processes) which in themselves may also be contributing an important role. There is therefore, at this stage, a pressing need for future research that not only evaluates the importance of cognitions in the successful outcome of CBT for PDA but which also takes into account the possible role of these other mechanisms in achieving such outcome.

REFERENCES

- American Psychiatric Association (A.P.A.) (1980). Diagnostic and statistical manual of mental disorders (3rd ed.). Washington, DC: Author.
- American Psychiatric Association (A.P.A.) (1987). Diagnostic and statistical manual of mental disorders (3rd ed. - revised). Washington, DC: Author.
- American Psychiatric Association (A.P.A.) (1994). Diagnostic and statistical manual of mental disorders (4th ed.). Washington, DC: Author.
- Arnkoff, D.B., & Glass, C.R. (1989). Cognitive assessment in social anxiety and social phobia. Clinical Psychology Review, *9*, 61-74.
- Ascher, L.M. (1981). Employing paradoxical intention in the treatment of agoraphobia. Behaviour Research and Therapy, *19*, 533-542.
- Ascher, L.M., Schotte, D.E., & Grayson, J.B. (1986). Enhancing effectiveness of paradoxical intention in treating travel restriction in agoraphobia. Behaviour Therapy, *17*, 124-130.
- Bandura, A. (1977). Self-efficacy: Towards a unifying theory of behavioural change. Psychological Review, *84*, 191-215.
- Bandura, A., Taylor, C.B., Williams, S.L., Mefford, I.N. & Beauchas, J.D. (1985). Catcholamine secretion as a function of perceived coping self efficacy. Journal of Consulting and Clinical Psychology, *53*, 406-414.
- Barlow, D.H., Mavissakalian, M., & Hay, L.R. (1981). Couples treatment of agoraphobia: changes in marital satisfaction. Behaviour Research and Therapy, *19*, 245-255.
- Barlow, D.H., O'Brien, G.T., & Last, C.G. (1984). Couples treatment of agoraphobia. Behaviour Therapy, *15*, 41-58.
- Beamish, P.M., Granello, P.F., Granello, D.H., McSteen, P.B., et al. (1996). Outcome studies in the treatment of panic disorder: A review. Journal of Counseling and Development, *74*, 460-467
- Beck G.J., Stanley, M.A., Baldwin, L.E., Deagle, E.A., Averill, P.M. (1994). Comparison of cognitive therapy and relaxation training for panic disorder. Journal of Consulting and Clinical Psychology, *62*, 818-826.
- Beck, A.T. (1988). Cognitive approaches to panic disorder: Theory and therapy. In S. Rachman & J.D. Maser (Eds.), Panic: Psychological Perspectives (pp. 91-110). New Jersey: Lawrence Erlbaum.
- Bouchard, S, Gauthier, J., Laberge, B., French, D., Pelletier, M., Godbout, C. (1996). Exposure versus cognitive restructuring in the treatment of panic disorder with agoraphobia Behaviour Research and Therapy, *34*, 213-224.
- Buttler, G. (1989). Issues in the application of cognitive and behavioural strategies to the treatment of social phobia. Clinical Psychology Review, *9*, 91-106.
- Chambless, D.L., Caputo, G.C., Bright, P., & Gallagher, R. (1984). The assessment of fear in agoraphobics: The body sensations questionnaire and the agoraphobic cognitions questionnaire. Journal of Consulting and Clinical Psychology, *52*, 1090-1097.
- Chambless, D.L. & Gillis, M.M. (1993). Cognitive therapy of anxiety disorders. Special Section: Recent developments in cognitive and constructivist psychotherapies. Journal of Consulting and Clinical Psychology, *61*, 248-260.
- Chambless, D.L., Goldstein, A.J., Gallagher, R., & Bright, P. (1986). Integrating behaviour therapy and psychotherapy in the treatment of agoraphobia. Psychotherapy, *23*, 150-159.
- Chambless, D.L., & Williams, K.E. (1995). A preliminary study of african americans with agoraphobia: Symptom severity and outcome of treatment with in vivo exposure. Behavior Therapy, *26*, 501-515.
- Clark, D.M. (1988). A cognitive model of panic attacks. In S. Rachman & J.D. Maser (Eds.), Panic: Psychological Perspectives (pp. 71-90). New Jersey: Lawrence Erlbaum.

de Beurs, E., Lange, A., van Dyck, R., Koele, P. (1995). Respiratory training prior to exposure in vivo in the treatment of panic disorder with agoraphobia: Efficacy and predictors of outcome. *Australian and New Zealand Journal of Psychiatry*, *29*, 104-113.

de Beurs, E., Lange, A., Koele, P., van Dyck, R.M.D., (1993) Frequency of panic as an outcome measure in agoraphobia research: Latent effects of exposure on panic. *Journal of Anxiety Disorders*, *7*, 307-319.

de Beurs, E., van Balkom, A.J.L.M., Lange, A., Koele, P., van Dyck, R.M.D. (1995) Treatment of panic disorder with agoraphobia: Comparison of fluvoxamine, placebo, and psychological panic management combined with exposure and of exposure in vivo alone *American Journal of psychiatry*, *152*, 683-691.

de Voge, J.T., Minor, T., & Karoly, P. (1981). Effects of behavioural intervention and interpersonal feedback on fear and avoidance components of severe agoraphobia: A case analysis. *Psychological Reports*, *49*, 595-605.

Emmelkamp, P.M.G., Kuipers, A.C. & Eggeraat, J.B. (1978). Cognitive modification versus prolonged exposure in vivo: A comparison with agoraphobics as subjects. *Behaviour Research and Therapy*, *16*, 33-41

Emmelkamp, P.M.G. (1982). *Phobic and obsessive-compulsive disorders*. New York: Guilford.

Emmelkamp, P.M.G., & Mersch, P.P. (1982). Cognitive and exposure in vivo in the treatment of agoraphobia: Short-term and delayed effects. *Cognitive Therapy and Research*, *6*, 77-88.

Fava, A.G., Zielezny, M., Savron, G., Grandi, S. (1995). Long term effects of behavioural treatment for panic disorder with agoraphobia *British Journal of Psychiatry*, *166*, 86-92.

Free, M.L., & Oei, T.P.S. (1989). Biological and psychological processes in the treatment and maintenance of depression. *Clinical Psychology Review*, *9*, 653-688.

Heimberg, R.G. (1989). Cognitive and behavioural treatments for social phobia: A critical analysis. *Clinical Psychology Review*, *9*, 187-198.

Himadi, W.G., Boice, R., & Barlow, D.H. (1985). Assessment of agoraphobia: Triple response measurement. *Behaviour Research and Therapy*, *23*, 311-323.

Hoffart, A. (1995). A comparison of cognitive and guided mastery therapy of agoraphobia *Behaviour Research and Therapy*, *33*, 423-434.

Jacobson, N.S., Follette, W.C., & Revensdorf, D. (1984). Psychotherapy outcome research: Methods for reporting variability and evaluating clinical significance. *Behavior Therapy*, *15*, 336-352.

Keijsers, G.P.J., Hoogduin, C.A.L., Schapp, C.P.D.R. (1994) Prognostic factors in the behavioral treatment of panic disorder with and without agoraphobia. *Behavior Therapy*, *25*, 689-708.

Khawaja, N.G. & Oei, T.P.S. (In press). Catastrophic cognitions in panic disorder with and without agoraphobia: A Review. *Clinical Psychology Review*.

Lang, P.J. (1988). Fear, anxiety, and panic: Context, cognition, and visceral arousal. In S. Rachman & J.D. Maser (Eds.), *Panic: Psychological Perspectives* (pp. 219-236). New Jersey: Lawrence Erlbaum.

Last, C.G., Barlow, D.H., & O'Brien, G.T. (1984). Cognitive change during treatment of agoraphobia: Behavioural and cognitive behavioural approaches. *Behaviour Modification*, *8*, 181-210.

Mackay, W., & Liddell, A. (1986). An investigation into the matching of specific agoraphobic anxiety response characteristics with specific types of treatment. *Behaviour Research and Therapy*, *24*, 361-364.

Marchione, K.E., Michelson, L., Greenwald, M., & Dancu, C. (1987). Cognitive behavioural treatment of agoraphobia. *Behaviour Research and Therapy*, *25*, 319-328.

Marks, I.M., Swinson, R.P., Basoglu, M., Kuch, K., Noshirvani, H., O'Sullivan, G., Lelliott, P.T., Kirby, M., McNamee, G., Sengun, S., Wickwire, Kim., (1993) Alprazolam and exposure alone and combined in panic disorder with agoraphobia: A controlled study in London and Toronto. *British Journal of Psychiatry*, *162*, 776-787.

- Mavissakalian, M., Michelson, L., Greenward, D., Kornblith, S., & Greenwald, M. (1983). Cognitive-behavioural treatment of agoraphobia: Paradoxical intention vs self-statement training. Behaviour Research and Therapy, *21*, 75-86.
- Michelson, L., Marchione, K., Greenwald, M., Glanz, L., Testa, S., & Marchione, N. (1990). Panic disorder: Cognitive-behavioral treatment. Behaviour Research and Therapy, *28*, 141-151.
- Michelson, L., Mavissakalian, M., & Marchione, K. (1985). Cognitive and behavioral treatments of agoraphobia: Clinical, behavioural, and psychophysiological outcomes. Journal of Consulting and Clinical Psychology, *53*, 913-925.
- Michelson, L., Mavissakalian, M., & Marchione, K. (1988). Cognitive, behavioural, and psychophysiological treatments of agoraphobia: A comparative outcome investigation. Behavior Therapy, *19*, 97-120.
- Michelson, L., Mavissakalian, M., Marchione, K., Dancu, C., & Greenwald, M. (1986). The role of self-directed in vivo exposure in cognitive, behavioural, and psychophysiological treatments of agoraphobia. Behavior Therapy, *17*, 91-108.
- Nietzel, M.T., & Trull, T.J., (1988). Meta-analytic approaches to social comparisons: A method for measuring clinical significance. Behavioral Assessment, *10*, 159-169.
- O'Brien, J.S. (1979). A modified thought stopping procedure for the treatment of agoraphobia. Journal of Behavior Therapy and Experimental Psychiatry, *10*, 121-124.
- Oei, T.P.S., Duckham, S., & Free, M. (1989). Does cognitive behaviour therapy support cognitive models of depression? Behaviour Change, *6*, 70-75.
- Oei, T.P.S., & Free, M.L. (1995). Do cognitive behaviour therapies validate cognitive models of mood disorders? A review of the empirical evidence. International Journal of Psychology, *30*, 145-180.
- Oei, T.P.S., Lim, B., & Young, R.M. (1991). Cognitive processes and cognitive behavior therapy in the treatment of problem drinking. Journal of Addictive Diseases, *10*, 63-80.
- Oei, T.P.S. & Shuttlewood, G.J. (1996). Specific and non specific factors in psychotherapy: A case of cognitive therapy for depression. Clinical Psychology Review, *16*, 83-103.
- Ollendick, T.H., (1995). Cognitive behavioral treatment of panic disorder with agoraphobia in adolescents: A multiple baseline design analysis. Behavior Therapy, *26*, 517-531.
- Ost, L-G., Westling, B.E., Hellstrom, K. (1993). Applied relaxation, exposure in vivo and cognitive methods in the treatment of panic disorder with agoraphobia. Behaviour Research and Therapy, *31*, 383-394.
- Powell, M.B. & Oie, T.P.S. (1991). Cognitive processes underlying the behavior change in cognitive behavior therapy with childhood disorders: A review of experimental evidence. Behavioural Psychotherapy, *19*, 247-265
- Rachman, S., & Maser, J.D. (Eds.) (1988). Panic: Psychological perspectives. New Jersey: Lawrence Erlbaum.
- Rapee, R. (1986). Differential response to hyperventilation in panic disorder and generalised anxiety disorders. Journal of Abnormal Psychology, *95*, 24-28.
- Rapee, R. (1987). The psychological treatment of panic attacks: Theoretical conceptualisation and review of evidence. Clinical Psychological Review, *7*, 427-438.
- Salkovskis, P.M. (1988). Phenomenology, assessment, and the cognitive model of panic. In S. Rachman & J.D. Maser (Eds.), Panic: Psychological Perspectives (pp. 111-136). New Jersey: Lawrence Erlbaum.
- Salkovskis, P.M. Clark, D.M., Hackmann, A. (1991) Treatment of panic attacks using cognitive therapy without exposure or breathing retraining. Behaviour Research and Therapy, *29*, 161-166.
- Salkovskis, P.M. Clark, D.M., Hackmann, A., Wells, A., Gelder, M.G. (In Submission). An experimental investigation of the role of safety behaviours in the maintenance of panic disorder with agoraphobia. Journal of Abnormal Psychology.

Salkovskis, P.M., Jones, D.R.O., & Clark, D.M. (1986). Respiratory control in the treatment of panic attacks: replication and extension with concurrent measures of behaviour and pCO₂. British Journal of Psychiatry, *148*, 526-532.

Sharp, D.M., Power, K.G., Simpson, R.J., Swanson, V., Moodie, E., Anstee J.A., Pharm, B., Ashford, J.J. (1996). Fluvoxamine, placebo, and cognitive behaviour therapy used alone and in combination in the treatment of panic disorder and agoraphobia. Journal of Anxiety Disorders, *10*, 219–242.

Swinson, R.P., Fergus, K.D., Cox, B.J. Wickwire, K. (1995) Efficacy of telephone-administered behavioural therapy for panic disorder with agoraphobia. Behaviour Research and Therapy, *33*, 465–469.

Taylor, S. Woody, S., Koch, W.J., McLean, P.D, Anderson, K.W. (1996). Suffocation false alarms and efficacy of cognitive behavioral therapy for panic disorder. Behavior Therapy, *27*, 115–126.

Trull, T.J., Nietzel, M.T., & Main, A. (1988). The use of meta-analysis to assess the clinical significance of behaviour therapy for agoraphobia. Behavior Therapy, *19*, 527-538.

Williams, S.L., & Rappoport, A. (1983). Cognitive treatment in the natural environment for agoraphobics. Behavior Therapy, *14*, 299-313.

Table 1

Summary of CBT Studies on Panic Disorder With Agoraphobia

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
1/ Ascher (1981)	9 Females 1 Male Group comparison Multiple baseline Sequential Staggering of treatments	1. CBT (GE + PI) 2. CBT (PI)	Behavioural	None	No	NA	CBT Effective 2 > 1
2/ Ascher, Schotte & Grayson (1986)	13 Females 2 Males Randomised Group Comparison	1. CBT (PI) 2. CBT (PI + EP) 3. CBT (EP)	Behavioural	None	No	NA	CBT Effective 2 > 1 > 3
3/ Barlow, Mavissakalian & Hay (1981)	6 Females & Husbands Group Comparison	1. CBT (CR + CC + GE) with husbands	Fear/Avoidance Marital State	6-16 months	No	NA	CBT Effective

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
4/ Barlow, O'Brien & Last (1984)	28 Females & Husbands Randomised Group Comparison	1. CBT (CST + PI + CR + GE) with husbands 2. CBT (CST + PI + CR + GE) without husbands	Panic Fear/Avoidance Severity/Intensity General Symptoms Behavioural Depression Marital State Composite	None	No	NA	CBT effective 1 > 2
5/ Beck, Stanley, Baldwin, Deagle, Averill (1994)	52 Female 18 Male Randomised Group Comparison	1. CBT (CR) 2. CBT (RT) 3. Minimal Contact Control	Panic Fear Severity / Intensity Anxiety Social Anxiety Cognitive Physiological Depression	1, 3 & 6 month	Yes	1. +/+ 2. +/+ 3. -	CBT effective. 1 & 2 > 3. Support for 1 > 2 on specific cognitive changes, although not strongly.

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
6/ Bouchard, Gauthier, Laberge, French, Pelletier & Godbout (1996)	24 Female 4 Male Random Group Comparison	1. CBT (Interceptive & exteroceptive exposure) 2. CBT (CR)	Panic Fear/Avoidance Anxiety Cognitive Self-Efficacy Depression Endstate Funct.	6 month	Yes	1. +/+ 2. +/+	CBT effective 1 & 2 produce change over time. 1=2
7/ Chambless, Goldstein, Gallagher & Bright (1986)	30 Females 5 Males Group Comparison Controls	1. CBT (PI + TS + CR + GT + YB + SR + FST + GE) 2.WLC	Panic Fear/Avoidance Anxiety Social Anxiety Cognitive Depression Marital State Assertiveness	6 month	Yes	1. +/ + 2. NA/ NA	CBT Effective 1 > 2
8/ Chambless & Williams (1995)	65 Females 10 Males African American Vs White (Post Hoc design)	Post Hoc with various CBT Treatments. All included GE	Panic Fear/Avoidance Severity/Intensity Behavioural Depression	6 month	Yes	+	CBT (in vivo exposure) effective. Black patients improved less on measures of phobia.

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
9/ DeBeurs, Lange, VanDyck, Koele (1995)	N=32 (sex not specified)	1. CBT (BT + GPE)	Panic Fear/Avoidance General Symptom Behavioural Depression	3 & 6 month	No	NA	CBT effective. Higher pre-treatment agrophobic complaints, use of psychotropic medication & longer duration of disorder associated with poorer outcome.
10/ DeBeurs, VanBalkom, Lange, Koele, VanDyck (1995)	N=96 (sex not specified) Randomised group comparison	1. Fluvoxamine + GPE 2. Placebo + GPE 3. CBT (BT + GPE) 4. GPE	Panic Fear/Avoidance Severity/Intensity General Symptom Depression Endstate Funct.	No	Yes	1. + 2. + 3. + 4. +	1, 2, 3, 4 effective. 1 > 2, 3, 4. 2, 3 & 4 equally effective.
11/ DeBeurs, Lange, Koele & VanDyck (1993)	N=28 (sex not specified)	1. CBT (BT + GPE)	Panic Fear/Avoidance Behavioural Endstate Funct.	None	No	NA	CBT Effective
12/ DeVoge, Minor & Karoly (1981)	1 Male Single case Sequential adding of treatments	1. CBT (RT + SST + CR + IF)	Severity/Intensity	16 month	Yes	1. +/ NR	CBT Effective

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
13/ Emmelkamp Kuipers & Eggeraat (1978)	20 Females & Males Randomised Cross-over Group Comparison	1. CBT (CR) 2. PE	Fear/Avoidance Anxiety Behavioural Locus of Control Depression Assertiveness	1 month	No	NA	1 & 2 Effective 2 > 1
14/ Emmelkamp & Mersch (1982)	22 Females 5 Males Group Comparison	1. PE 2. CBT (CR) 3. CBT (SST + PE)	Fear/Avoidance Anxiety Behavioural Locus of Control Depression Assertiveness	1 month	No	NA	1, 2 & 3 Effective 1 & 3 > 2 at post- treatment 1 = 2 = 3 at F/Up
15/ Fava, Zielezny, Savron & Grandi (1995)	N=93 (sex not specified)	1. CBT (exposure in vivo only)	Panic Fear/Avoidance	2-9 years	No	NA	81 became panic free by post -treatment. Continued remission rates = 96.1% for >2years, 77.6% >4years and 67.4% >7 years.

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
16/ Hoffart (1995)	N=52 (sex not specified) Rondomised Group Comparison	1. CBT (CT) 2. CBT (Guided Mastery)	Panic Fear/Avoidance Severity/Intensity Anxiety Social Anxiety Behavioural Cognitive Self-Efficacy Depression Endstate Funct.	No	Yes	1. + 2. +	CBT effective. 1 produced greater change on ACQ than 2. Sig. more patients in 1 attained a high endstate functioning on fears. However, while no overall sig difference between 1 & 2 on continuous outcome measures, analyses showed CT to improve more consistently across measures.
17/ Keijsers, Hoogduin & Schaap (1994)	44 Females 16 Males	1. CBT (SIT + GE)	Panic Fear/Avoidance Anxiety Cognitive Depression Marital State	2 month	Yes	+ / +	CBT effective. Catastrophic cognitions best predictor of poor outcome. Patients affected by maladaptive cognitions improved less with behavioural intervention.

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
18/ Last, Barlow & O'Brien (1984)	5 Females 1 Male Matched to severity of disorder Multiple baseline	1. GE 2. CBT (SST + GE)	Fear/Avoidance Severity/Intensity Behavioural Cognitive Physiological	1 month	Yes	1. Variable/ Variable 2. Variable/ Variable	Treatment effectiveness variable across patients
19/ Mackay & Liddell (1986)	11 Females 3 Males Half of the patients matched for mode of responding to treatment	1. CBT (CT + E) matched group 2. CBT (CT + E) non-matched group 3. RT + E matched group 4. RT + E non-matched group	General Symptoms Cognitive	6 month	Yes	NR/ +	1, 2, 3 & 4 effective. 1 & 4 > 2 & 3; 1 > 2; 4 > 3 post-treatment 3 & 4 > 1 & 2 at F/Up

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
20/ Marchione, Michelson, Greenwald & Dancu (1987)	14 Females Randomised Group Comparison	1. GE 2. RT + GE 3. CBT (CT + GE)	Panic Fear/Avoidance Severity/Intensity Anxiety General Symptoms Behavioural Cognitive Physiological Depression Composite	No	Yes	1. NS/ NR 2. +/ NR 3. +/ NR	1, 2 & 3 effective 2 & 3 > 1
21/ Marks, Swinson, Basoglu, Kuch, Noshirvani, O'Sullivan, Lelliott, Kirby, McNamee, Sengun, Wickwire (1993)	117 Females 27 Males Randomised	1. Alprazolam + Guided exposure 2. Alprazolam + Relaxation 3. Placebo + Guided Exposure 4. Placebo + Relaxation	Panic Fear/Avoidance Severity/Intensity Anxiety Depression	6 month	No	NA	1, 2, 3, & 4 all improved throughout. 1 improved gains during tx, but impaired improvement thereafter. Exposure had twice the effect size of alprazolam and tx gains maintained at follow-up, unlike alprazolam. Relapse likely after alprazolam stopped.

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
22/ Mavissak- alian, Michelson, Greenwald, Kornblith & Greenwald (1983)	24 Females & Males Randomised Group Comparison	1. CBT (SST + E) 2. CBT (PI + E)	Fear/Avoidance Severity/Intensity Anxiety Behavioural Cognitive Depression	1 & 6 month	Yes	1. +/- NR 2. +/- NR	1 & 2 effective 2 > 1 at post-treatment; 2 = 1 at F/Up
23/ Michelson, Mavissak- alian, & Marchione (1985)	27 Females 4 Males Randomised Group Comparison	1. GE 2. CBT (PI + SDP) 3. RT + SDP	Panic Fear/Avoidance Severity/Intensity Anxiety Behavioural Physiological Locus of Control Depression Assertiveness Composite	3 month	No	NA	1, 2 & 3 effective 1 > 3 > 2

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
24/ Michelson, Mavissak- alian, Marchione, Dancu & Greenwald (1986)	27 Females 4 Males Randomised Group Comparison	1. GE 2. CBT (PI + SDP) 3. RT + SDP	Fear/Avoidance Severity/Intensity Behavioural Depression Composite	3 month	No	NA	1, 2,& 3 effective
25/ Michelson, Mavissak- alian & Marchione (1988)	57 Females 16 Males Randomised Group Comparison	1. GE 2. CBT (PI + SDP) 3. RT + SDP	Panic Fear/Avoidance Severity/Intensity Anxiety General Symptoms Behavioural Locus of Control Depression Marital State Assertiveness Composite	3 month	No	NA	1, 2 & 3 effective 1 = 2 = 3

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
26/ Michelson, Marchione, Greenwald, Glanz, Testa, Marchione (1990)	6 Females 4 Males	1. CBT (CT + RT)	Panic Fear/Avoidance Severity/Intensity Anxiety General Symptom Cognitive Physiological Depression Marital State	No	Yes	+	CBT effective. All S's panic free by post-treatment.
27/ O'Brien (1979)	1 Female 1 Male Single cases	1. CBT (TS)	Severity/Intensity	12 month	No	NA	CBT effective
28/ Ollendick (1995)	3 Female adolescents 1 Male adolescent Multiple base line single cases design	1. CBT (SIT + GE + CT)	Panic Fear/Avoidance Anxiety Self-Efficacy Depression	6 month	No	NA	CBT effective with adolescents with PDA.
29/ Ost, Westling & Hellstrom (1993)	30 Female 15 Male Randomised	1. GE 2. RT 3. CT (self-talk + attribution training +CR)	Fear/Avoidance Anxiety Behavioural Cognitive Depression	12 month	Yes	1. +/+ 2. +/+ 3. +/+	CBT effective. 1, 2 & 3 all improved by post tx and maintained improvement at follow-up.

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
30/ Salkovskis Clark & Hackmann (1991)	N=7 (sex not specified) Multiple baseline	1. Focal CT 2. Non-Focal CT	Panic Cognitive	4 week	Yes	1. +/+ 2. -/NA	Cognitive techniques aimed at misinterpretations of bodily sensations reduce panic, yet cognitive procedures not targeting this don't.
31/ Salkovskis Jones & Clark (1986)	9 Females 1 Male Matched Group	1. CBT (RASS + BT)	Panic Depression	6 month	No	NA	CBT effective
32/ Sharp et al. (1996)	115 Female 32 Male Randomised Group Comparison	1. Fluvoxamine 2. Placebo 3. F + CBT 4. P + CBT 5. CBT	Panic Fear/Avoidance Severity/Intensity Anxiety Depression	6 month	No	NA	CBT effective. All active tx groups improved. CBT groups preserved tx gains better over time. F + CBT improved quicker.
33/ Swinson, Fergus, Cox, Wickwire (1995)	37 Female 5 Male Randomised	1. Telephone CBT (GE) 2. WLC	Fear/Avoidance Severity/Intensity Anxiety General Symptom Depression	3 month & 6 month	No	NA	CBT effective. Exposure based treatment (1) improved (& maintained at f/up). Wait list did not improve and then once treated with 1 improved.

Study Number / Author	Design & Subjects	Treatment Groups	Outcome Measures	F/Up	Cognitive Measure Used ?	Change In Cognitive Measure Post/ F-Up	Outcome and Comments
34/ Taylor, Woody, Koch, McLean, Anderson (1996)	16 Female 6 Male Group design by suffocation panickers or non suffocation panickers (PD criteria the only necessity).	1. CBT	Panic Fear/Avoidance Anxiety	3 Month	No	NA	CBT effective at post-treatment & follow-up for both suffocation and nonsuffocation panickers.
35/ Williams & Rappoport (1983)	20 Females Randomised Group Comparison	1. CBT (CTT + GE) 2. GE	Anxiety Behavioural Cognitive	3 to 5 months	Yes	1. +/- NR 2. +/- NR	CBT effective

Key: ACQ=agoraphobic cognitions questionnaire; BT=breathing training; CC=covert rehearsal of coping with anxiety; CR=cognitive restructuring; CT=cognitive therapy; CTT=coping thoughts training; CST=coping self-statement training; E=exposure; EP=enhancement procedures; F=fluvoxamine; FST=family systems therapy; F-Up=follow-up; GE=graduated exposure; GPE=graduated prolonged exposure; GT=gestalt therapy; IF=interpersonal feedback; NA=not applicable; NR=not reported; OESF/I=operationalised measures of endstate functioning and improvement; P=placebo; PE=prolonged exposure; PI=paradoxical intervention; RASS=retribution of somatic symptoms; RT=relaxation training; SDP=self-directed practice; SIT=stress inoculation training; SR=self-reinforcement; SUSs=subjective units of discomfort scale; TS=thought stopping; WLC=wait list control; YB=yogic breathing.

Table 2

Statistical Improvement, Within Domains, by Study

Measures of:	Improved by Post-tx &/ or Follow-up	No Improvement
Panic	4, 5, 6, 7, 8, 9, 10, 11, 15, 16, 17, 20, 21, 23, 25, 26, 28, 30, 31, 32, 34.	
Fear & Avoidance	3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26, 28, 29, 32, 33, 34.	
Severity / Intensity	5, 8, 10, 12, 16, 18, 20, 21, 22, 23, 24, 25, 26, 27, 32, 33.	
General Anxiety	5, 6, 7, 14, 16, 17, 21, 22, 23, 25, 26, 28, 29, 32, 33, 34, 35.	13, 20.
Social Anxiety	7, 16, 26, 29.	5.
General Symptomatology	4, 5, 9, 10, 19, 20, 25, 26, 33.	20.
Behavioural	1, 2, 8, 9, 10, 11, 13, 14, 16, (18), 20, 23, 25, 29, 35.	22.
Cognitive	5, 6, 7, 8, 10, 16, 17, 26, 29, 20, 22, 26, 29, 30, 35.	18.
Physiological	5, 18, 20, 23, 26.	5, 18.
Locus of Control	6, 13, 14, 16, 23, 25, 28.	
Depression	4, 5, 6, 7, 8, 9, 10, 13, 14, 16, 17, 20, 21, 22, 23, 24, 25, 26, 28, 29, 31, 32, 33.	
Marital State	3, 17, 25, 26.	3.
Assertiveness	7, 14, 23, 25.	13.
Endstate Functioning	4, 5, 6, 10, 11, 16, 20, 23, 24, 25.	

Note: Where a study found significant reduction within a domain on one measure and not on another, it has been entered in both categories of improved and not improved. Any study in brackets () signifies mixed results.

Table 3

Pre, Post and Follow-up Means and Effect Size Scores for the FQ Agoraphobia Subscale Scores for Studies in the Meta-Analysis

Study	Pre-Test Mean	Effect Size 1	Effect Size 2	Post-Test Mean	Effect Size 1	Effect Size 2	F/Up Mean	Effect Size 1	Effect Size 2
Barlow et al. (1984)	27.00	4.60	1.76	17.90	2.67	0.70	----	----	----
Beck et al. (1994)	1). 13.07 2). 11.50 3). 9.77	1.65 1.32 0.95	0.15 -0.03 -0.24	5.73 10.06 9.62	0.09 1.01 0.92	-0.71 -0.21 -0.26	----	----	----
Chambless et al. (1986)	26.39	4.47	1.69	12.09	1.44	0.03	10.09	1.01	0.56
DeBeurs et al. (1995)	32.90	5.87	2.45	22.20	3.74	1.29	----	----	----
Marchione et al. (1987)	29.80	5.19	2.09	10.6	1.12	-0.14	----	----	----
Mavissakalian et al. (1983)	1). 22.0 2). 23.0	3.55 3.77	1.18 1.30	17.7 13.6	2.64 1.77	0.68 0.21	12.30 13.00	1.49 1.64	0.05 0.14
Michelson et al. (1988)	28.0	4.83	1.88	8.60	0.70	-0.38	10.90	1.19	-0.11
Michelson et al. (1990)	12.11	1.43	0.03	3.56	-0.38	-0.96	----	----	----
Ost et al. (1993)	1). 25.20 2). 27.27 3). 26.80	4.23 4.67 4.57	1.56 1.80 1.74	12.60 11.53 15.80	1.55 1.33 2.23	0.09 -0.03 0.47	11.80 12.40 14.60	1.38 1.78 1.98	0.00 0.07 0.33
Swinson et al. (1995)	1). 25.50 2). 27.65	4.30 4.76	1.59 1.84	17.30 27.18	2.55 4.66	0.64 1.79	} 15.69	2.19	0.45
Williams & Rappoport (1983)	1). 22.60 2). 24.10	3.68 3.97	1.25 1.43	16.80 16.00	2.45 2.25	0.58 0.48	15.80 15.40	2.23 2.13	0.46 0.42

Note: Effect Size 1 is the means of the studies compared with Trull and colleagues (1988) college norms for the FQ agoraphobia subscale, and Effect Size 2 is the means of the studies compared to the general (community) population norms for the FQ agoraphobia subscale.

Table 4

Pre, Post and Follow-up Means and Effect Size Scores for the FQ Total Scale Scores for Studies in the Meta-Analysis


Study	Pre-Test Mean	Effect Size 1	Effect Size 2	Post-Test Mean	Effect Size 1	Effect Size 2	F/Up	Effect Size 1	Effect Size 2
Marchione et al. (1987)	67.60	3.01	1.67	22.20	-0.42	-1.12	----	----	----
Mavissakalia et al. (1983)	1). 51.9 2). 47.4	1.83 1.49	0.71 0.43	45.1 29.1	1.31 0.10	0.29 -0.70	36.9 27.9	0.69 0.01	-0.22 -0.77
Michelson et al. (1988)	59.50	2.40	1.17	26.10	-0.12	-0.88	29.30	0.12	-0.69
Michelson et al. (1990)	63.00	2.66	1.39	32.78	0.38	-0.47	----	----	----
Ost et al. (1993)	1). 52.27 2). 52.07 3). 51.47	1.85 1.84 1.79	0.73 0.72 0.68	32.87 28.40 32.33	0.39 0.05 0.35	-0.47 -0.74 -0.50	30.00 27.80 28.47	0.22 0.01 0.06	-0.64 -0.78 -0.74
Salkovskis et al. (1986)	53.20	1.92	1.17	28.44	0.06	-0.74	25.11	-0.20	-0.16
Williams & Rappoport (1983)	1). 56.10 2). 57.80	2.14 2.27	0.96 1.07	41.00 41.40	1.00 1.03	0.03 0.06	41.01 36.60	1.01 0.67	0.04 -0.24

Note: Effect Size 1 is the means of the studies compared with Trull and colleagues (1988) college norms for the FQ total scale, and Effect Size 2 is the means of the studies compared to the general (community) population norms for the FQ total scale.


FIGURE CAPTIONS

Figure 1. Mean effect sizes for the agoraphobia subscale of the fear questionnaire plotted at pre and post-treatment and follow-up.

Figure 2. Mean effect sizes for the total phobia subscale of the fear questionnaire plotted at pre and post-treatment and follow-up.



■ College Norms
● General Norms



- College Norms
- General Norms